arXiv:1109.0715v2 [math.QA] 16 Oct 2011

Connection Problem of Knizhnik-Zamolodchikov Equation on Moduli Space $\mathcal{M}_{0,5}$

OI, Shu[*] UENO, Kimio[†]

Abstract

In this article, we consider the connection problem of the KZ (Knizhnik-Zamolodchikov) equation on the moduli space $\mathcal{M}_{0,5}$, and show that the connection matrices are expressed in terms of the Drinfel'd associator. As the compatibility condition on the connection problem, we have the pentagon relation for the Drinfeld associators. As an application of the connection problem, we derive the five term relation for dilogarithms.

1 Introduction

In this article, we consider the connection problem of the KZ equation on the moduli space $\mathcal{M}_{0,5}$, and show that the connection matrices are expressed in terms of the Drinfel'd associators. As the compatibility condition on the connection problem, we have the pentagon relation for the Drinfeld associators. This story was already proved by Drinfel'd in his famous paper [Dr] (see also Wojtkowiak's paper [W]), but our proof is completely different from theirs. Our proof is based upon the decomposition theorem for fundamental solutions of the KZ equation. As an application of the connection problem, we derive the five term relation for dilogarithms.

This paper is organized as follows: In section 2, we introduce the KZ equation on the moduli space $\mathcal{M}_{0,n}$. In the cubic coordinate system of the space, it is represented as the KZ equation of $n-3$ variables. In section 3, we particularly consider the KZ equation of one variable which is denoted by z, and show that the Drinfel'd associator Φ_{KZ} gives the connection matrix between $\mathcal{L}(z)$, which is the fundamental solution of normalized at $z=0$ of the equation, and $\mathcal{L}^{(1)}(z)$,

---

[*] Department of Mathematics, School of Fundamental Sciences and Engineering, Faculty of Science and Engineering, Waseda University.
3-4-1, Okubo, Shinjuku-ku, Tokyo 169-8555, Japan.
e-mail: `shu_oi@toki.waseda.jp`

[†] Department of Mathematics, School of Fundamental Sciences and Engineering, Faculty of Science and Engineering, Waseda University.
3-4-1, Okubo, Shinjuku-ku, Tokyo 169-8555, Japan.
e-mail: `uenoki@waseda.jp`

which is the fundamental solution normalized at $z = 1$ of the equation;

$$\mathcal{L}(z) = \mathcal{L}^{(1)}(z)\mathit{\Phi}_{\mathrm{KZ}}. \tag{1}$$

In section 4, the KZ equation of two variables, which are denoted by (z_1, z_2), will be considered. The fundamental solution $\mathcal{L}(z_1, z_2)$ normalized at $(z_1, z_2) = (0, 0)$ of the equation is shown to decompose as follows:

$$\begin{aligned}\mathcal{L}(z_1, z_2) &= \mathcal{L}^{(2)}_{2\otimes 1}(z_1, z_2)\mathcal{L}^{(1)}_{2\otimes 1}(z_1) \\ &= \mathcal{L}^{(1)}_{1\otimes 2}(z_1, z_2)\mathcal{L}^{(2)}_{1\otimes 2}(z_2)\end{aligned} \tag{2}$$

where $\mathcal{L}^{(j)}_{i\otimes j}(z_j)$ is the fundamental solution normalized at $z_j = 0$ of the KZ equation of the one variable z_j, and $\mathcal{L}^{(j)}_{i\otimes j}(z_1, z_2)$ is the fundamental solution normalized at $z_i = 0$ of the Schlesinger type equation of the variable z_i. We will refer to (2) as "the decomposition theorem". In section 5, we will consider the connection problem of solutions of the KZ equation on $\mathcal{M}_{0,5}$. We fix a pentagon in $\overline{\mathcal{M}}_{0,5}(\mathbf{R})$, which denotes the real points of a certain compactification $\overline{\mathcal{M}}_{0,5}$ of $\mathcal{M}_{0,5}$, as below:



To each vertex P_α, we associate the cubic coordinate $(z_1^{(\alpha)}, z_2^{(\alpha)})$, and the KZ equation of the two variable $(z_1^{(\alpha)}, z_2^{(\alpha)})$. Let $\mathcal{L}^{(\alpha)}$ be the fundamental solution normalized at the vertex P_α of the equation. Then one can show that

$$\mathcal{L}^{(\alpha)} = \mathcal{L}^{(\alpha+1)}\mathit{\Phi}^{(\alpha)}_{\mathrm{KZ}} \quad (\alpha = 0, 1, 2, 3, 4 \pmod 5) \tag{3}$$

where $\mathit{\Phi}^{(\alpha)}_{\mathrm{KZ}}$ is the Drinfel'd associator attached to the vertex P_α. The relation (3) is proved by applying the decomposition theorem (2). As the compatibility condition for (3), we obtain the pentagon relation of the Drinfel'd associators

$$\mathit{\Phi}^{(4)}_{\mathrm{KZ}}\,\mathit{\Phi}^{(3)}_{\mathrm{KZ}}\,\mathit{\Phi}^{(2)}_{\mathrm{KZ}}\,\mathit{\Phi}^{(1)}_{\mathrm{KZ}}\,\mathit{\Phi}^{(0)}_{\mathrm{KZ}} = \mathbf{I} \tag{4}$$

In section 6, as an application of the connection problem, we derive the five term relation for dilogarithms

$$\begin{aligned}\mathrm{Li}_2(z_1 z_2) = \mathrm{Li}_2\left(\frac{-z_1(1-z_2)}{1-z_1}\right) + \mathrm{Li}_2\left(\frac{-z_2(1-z_1)}{1-z_2}\right) \\ + \mathrm{Li}_2(z_1) + \mathrm{Li}_2(z_2) + \frac{1}{2}\log^2\left(\frac{1-z_1}{1-z_2}\right).\end{aligned} \tag{5}$$

In [OU], we show that the decomposition theorem (2) is equivalent to the generalized harmonic product relations of hyperlogarithms of the type $\mathcal{M}_{0,5}$, which contain the harmonic product of multiple polylogarithms.

Acknowledgment The first author is partially supported by Waseda University Grant for Special Research Projects No. 2010B-200, 2011B-095. The second author is partially supported by JSPS Grant-in-Aid No. 19540056, 22540035.

2 The KZ equation on the moduli space $\mathcal{M}_{0,n}$

2.1 Definition of the moduli space and the KZ equation

The moduli space $\mathcal{M}_{0,n}$ is by definition [Y]

$$\mathcal{M}_{0,n} = \mathrm{PGL}(2, \mathbf{C}) \Big\backslash \mathbf{F}_n(\mathbf{P}^1). \tag{6}$$

Here $\mathbf{F}_n(\mathbf{P}^1)$ is the configuration space of n points on $\mathbf{P}^1 = \mathbf{P}^1(\mathbf{C})$,

$$\begin{aligned}\mathbf{F}_n(\mathbf{P}^1) &= \{(x_1, \dots, x_n) \in (\mathbf{P}^1)^n \mid x_i \neq x_j \ \ (i \neq j)\} \\ &= (\mathbf{P}^1)^n \setminus \Delta\end{aligned}$$

where Δ is the thick diagonal of $(\mathbf{P}^1)^n$,

$$\Delta = \{(x_1, \dots, x_n) \in (\mathbf{P}^1)^n \mid \exists i \neq j \text{ s.t. } x_i = x_j\}.$$

The KZ equation(KZE) on $\mathcal{M}_{0,n}$ is a Pfaffian system defined by

$$dG = \Omega G, \quad \Omega = \sum_{1 \le i < j \le n} \omega_{ij} \Omega_{ij}, \quad \omega_{ij} = d \log(x_i - x_j). \tag{KZE}$$

Here $\{\Omega_{ij}\}$ are generators of the Lie algebra $\mathfrak{X}$ of the fundamental group of $\pi_1(\mathcal{M}_{0,n})$,

$$\mathfrak{X} = \mathrm{Lie}(\pi_1(\mathcal{M}_{0,n})).$$

This Lie algebra is defined through the lower central series of $\pi_1(\mathcal{M}_{0,n})$,

$$\pi_1(\mathcal{M}_{0,n}) = G_1 \supset G_2 \supset G_3 \supset \cdots .$$

We set

$$\mathfrak{X} = \bigoplus_{j=1}^{\infty} \mathbf{C}\, G_j / G_{j+1}.$$

We should note that the fundamental group $\pi_1(\mathbf{F}_n(\mathbf{P}^1))$ of the configuration space $\mathbf{F}_n(\mathbf{P}^1)$ is isomorphic to

$$\pi_1(\mathbf{F}_n(\mathbf{P}^1)) \cong \pi_1(\mathcal{M}_{0,n}) \times \mathbf{Z}/2\mathbf{Z}$$

so that $\mathrm{Lie}(\pi_1(\mathbf{F}_n(\mathbf{P}^1)))$ is isomorphic to $\mathrm{Lie}(\pi_1(\mathcal{M}_{0,n}))$.

The generators $\{\Omega_{ij}\}_{1 \le i,j \le n}$ of $\mathfrak{X}$ satisfy the following relations [I]:

$$\Omega_{ij} = \Omega_{ji}, \quad \Omega_{ii} = 0, \tag{IH1}$$

$$\sum_{j=1}^{n} \Omega_{ij} = 0 \quad (\forall i), \tag{IH2}$$

$$[\Omega_{ij}, \Omega_{kl}] = 0 \quad (\{i, j\} \cap \{k, l\} = \emptyset). \tag{IH3}$$

These relations are referred to as the Ihara relations. $\mathfrak{X}$ is a Lie algebra generated by $\{\Omega_{ij}\}_{1\leq i,j\leq n}$ with the fundamental relations (IH1) $\sim$ (IH3). The relation (IH1) is rather convention of notations. From (IH2) and (IH3), we have

$$[\Omega_{ij}+\Omega_{jk},\Omega_{ik}]=0 \quad (\#\{i,j,k\}=3). \tag{7}$$

The one forms $\{\omega_{ij}\}_{1\leq i<j\leq n}$ satisfy the Arnold relations [A]:

$$\omega_{ij}\wedge\omega_{ik}+\omega_{ik}\wedge\omega_{jk}+\omega_{jk}\wedge\omega_{ij}=0 \tag{AR}$$

for $i<j<k$. It is known that (AR) are unique nontrivial relations of degree 2 which hold among these one forms.

From the Ihara relations (7) and the Arnold relations (AR), we can deduce that

(a) the system (KZE) is regular as $x_i=\infty$, $|x_j|<\infty$ $(j\neq i)$.

(b) the system (KZE) is integrable, namely Ω satisfy

$$\Omega\wedge\Omega=0.$$

(c) the system (KZE) is $\mathrm{PGL}(2,\mathbf{C})$ invariant.

Hence the system (KZE) can be regarded as an integrable system on $\mathcal{M}_{0,n}$.

2.2 Cubic coordinate system on $\mathcal{M}_{0,n}$ and KZ equation of $n-3$ variables

Let $r(i,j;k,l)=\dfrac{(x_i-x_k)(x_j-x_l)}{(x_i-x_l)(x_j-x_k)}$ be a cross ratio. Put

$$y_i=r(i,n-1;n-2,n)=\frac{x_i-x_{n-2}}{x_i-x_n}\frac{x_{n-1}-x_n}{x_{n-1}-x_{n-2}}\in\mathbf{P}^1\setminus\{0,1,\infty\} \tag{8}$$

for $i=1,\dots,n-3$. This defines a coordinate system which we call the simplicial coordinate system on $\mathcal{M}_{0,n}$. Now we introduce $(z_1,\dots,z_{n-3})$ by

$$y_i=z_1\cdots z_i \quad (1\leq i\leq n-3) \tag{9}$$

$$\Longleftrightarrow\begin{cases} z_1=y_1=r(1,n-1;n-2,n),\\ z_2=\dfrac{y_2}{y_1}=r(2,1;n-2,n),\\ \vdots\\ z_{n-3}=\dfrac{y_{n-3}}{y_{n-4}}=r(n-3,n-4;n-2,n).\end{cases} \tag{10}$$

This is also a coordinate system on $\mathcal{M}_{0,n}$ which we call the cubic coordinate system. These two coordinate system were introduced by F. Brown in [B].

In the cubic coordinate system on $\mathcal{M}_{0,n}$, (KZE) is represented as follows:

$$dG=\Omega G,$$
$$\Omega=\sum_{k=1}^{n-3}\xi_kX_k+\sum_{k=1}^{n-3}\xi_{kk}X_{kk}+\sum_{1\leq i<j\leq n-3}\xi_{ij}X_{ij}, \tag{11}$$

where

$$\xi_k = \frac{dz_k}{z_k}, \quad \xi_{kk} = \frac{dz_k}{1-z_k}, \quad \xi_{ij} = \frac{d(z_i \cdots z_j)}{1-(z_i \cdots z_j)} \quad (i<j) \tag{12}$$

and

$$X_k = \sum_{k \le i < j \le n-2} \Omega_{ij} \quad (1 \le k \le n-3), \quad X_{11} = -\Omega_{1,n-1}, \quad X_{kk} = -\Omega_{k-1,k},$$
$$X_{1j} = -\Omega_{j,n-1} \quad (2 \le j \le n-3), \quad X_{ij} = -\Omega_{i-1,j} \quad (2 \le i < j \le n-3).$$

We refer to this system as the KZ equation of $n-3$ variables.

3 KZE of one variable

3.1 Definition of KZE of one variable

By the cubic coordinate system $z = r(1,3;2,4) = \dfrac{(x_1-x_2)(x_3-x_4)}{(x_1-x_4)(x_3-x_2)}$, the moduli space $\mathcal{M}_{0,4}$ is homeomorphic to $\mathbf{P}^1 \setminus \{0,1,\infty\}$;

$$\mathcal{M}_{0,4} \xrightarrow{\sim} \mathbf{P}^1 \setminus \{0,1,\infty\} \ : \ [x_1,x_2,x_3,x_4] \mapsto z$$

where $[x_1,\ldots,x_4]$ stands for the $\mathrm{PGL}(2,\mathbf{C})$ orbit of $(x_1,\ldots,x_4)$. The KZ equation on $\mathcal{M}_{0,4}$ is represented as

$$\frac{dG}{dz} = \left(\frac{X_0}{z} + \frac{X_1}{1-z}\right) G \tag{KZE1}$$

where $X_0 = \Omega_{12}$, $X_1 = -\Omega_{13}$. This is the KZ equation of one variable.

In this case, $\mathfrak{X}$ is a free Lie algebra generated by X_0, X_1, which we denote by

$$\mathfrak{X} = \mathbf{C}\{X_0, X_1\}, \tag{13}$$

and the universal enveloping algebra $\mathcal{U}(\mathfrak{X})$ is a non-commutative polynomial algebra of X_0, X_1;

$$\mathcal{U} = \mathcal{U}(\mathfrak{X}) = \mathbf{C}\langle X_0, X_1\rangle. \tag{14}$$

3.2 Shuffle algebra of one forms

Put

$$\xi_0 = \frac{dz}{z}, \quad \xi_1 = \frac{dz}{1-z}. \tag{15}$$

They satisfy the Arnold relation $\xi_0 \wedge \xi_1 = 0$. We set

$$S = S(\xi_0,\xi_1) = \big(\mathbf{C}\langle \xi_0,\xi_1\rangle, ш, \mathbf{1}\big) \tag{16}$$

where $\mathbf{1}$ stands for the empty word in S and ш denotes the shuffle product, which is recursively defined by

$$w \,ш\, \mathbf{1} = \mathbf{1} \,ш\, w = w, \tag{17}$$
$$(\xi_{i_1} w_1) \,ш\, (\xi_{i_2} w_2) = \xi_{i_1}(w_1 \,ш\, (\xi_{i_2} w_2)) + \xi_{i_2}((\xi_{i_1} w_1) \,ш\, w_2). \tag{18}$$

S is an associative commutative algebra with the unit elemnt $\mathbf{1}$. This is called a shuffle algebra generated by ξ_0, ξ_1.

$\mathcal{U}$ is a non-commutative and cocomutative Hopf algebra, and S is a commutative Hopf algebra. As Hopf algebras, $\mathcal{U}$ and S are dual to each other.

To a word $w = \xi_{i_1} \cdots \xi_{i_r} \in S$, we can associate an iterated integral $\displaystyle\int_a^b w$ as follows [C1, Ha]:

$$\int_a^b \mathbf{1} = 1, \qquad \int_a^b w = \int_a^b \xi_{i_1}(t) \int_a^t \xi_{i_2} \cdots \xi_{i_r}. \tag{19}$$

Then we have

$$\int_a^b w_1 \int_a^b w_2 = \int_a^b (w_1 ш w_2). \tag{20}$$

For $w = \xi_0^{k_1-1}\xi_1 \cdots \xi_0^{k_r-1}\xi_1$, we can put an iterated integral $\displaystyle\int_0^z w$, which defines the multiple polylogarithm of one variable (MPL):

$$\mathrm{Li}_{k_1,\ldots,k_r}(z) := \int_0^z \xi_0^{k_1-1}\xi_1 \cdots \xi_0^{k_r-1}\xi_1. \tag{21}$$

This is a many-valued function on $\mathcal{M}_{0,4}$ and has a Taylor expansion

$$\mathrm{Li}_{k_1,\ldots,k_r}(z) = \sum_{n_1 > n_2 \cdots > n_r > 0} \frac{z^{n_1}}{n_1^{k_1} \cdots n_r^{k_r}} \quad (|z| < 1). \tag{22}$$

In particular if $k_1 \geq 2$, we can define an iterated integral $\displaystyle\int_0^1 \xi_0^{k_1-1}\xi_1 \cdots \xi_0^{k_r-1}\xi_1$. This is the multiple zeta value (MZV):

$$\zeta(k_1, \ldots, k_r) = \int_0^1 \xi_0^{k_1-1}\xi_1 \cdots \xi_0^{k_r-1}\xi_1 = \sum_{n_1 > n_2 \cdots > n_r > 0} \frac{1}{n_1^{k_1} \cdots n_r^{k_r}}. \tag{23}$$

Let us introduce shuffle subalgebras such as

$$S^0 = \mathbf{C1} \oplus S\xi_1, \quad S^{10} = \mathbf{C1} \oplus \xi_0 S \xi_1. \tag{24}$$

S^0 is a subalgebra spanned by words ending with other than ξ_0, and S^{10} is a subalgebra spanned by words beginning with other than ξ_1 and ending with other than ξ_0.

Proposition 1 ([R])**.** We have

$$S = S^0[\xi_0] = S^{10}[\xi_1, \xi_0]. \tag{25}$$

Let us define the regularization maps as follows:

$$\mathrm{reg}^0 : S = S^0[\xi_0] \to S^0, \quad u = \sum w_j ш \xi_0^j \mapsto w_0 \ (w_j \in S^0) \tag{26}$$

$$\mathrm{reg}^{10} : S = S^{10}[\xi_1, \xi_0] \to S^{10}, \quad u = \sum \xi_1^i ш w_{ij} ш \xi_0^j \mapsto w_{00} \ (w_{ij} \in S^{10}). \tag{27}$$

From Proposition 1, these maps turn out to be well-defined and homomorphisms. Furthermore we can also define the MPL map and the MZV map as follows:

- MPL map $\mathrm{Li}(\bullet;z):S\to\mathbf{C}$: For $u=\sum w_j ⧢ \xi_0^j \quad (w_j\in S^0)$, we put

$$\mathrm{Li}(u;z):=\sum \mathrm{Li}(w_j;z)\frac{\log^j z}{j!}. \tag{28}$$

  Here for a word $w=\xi_0^{k_1-1}\xi_1\cdots\xi_0^{k_r-1}\xi_1\in S^0$, $\mathrm{Li}(w;z)$ is the MPL $\mathrm{Li}_{k_1,\dots,k_r}(z)$.

- MZV map $\zeta:S^{10}\to\mathbf{C}$: For $w=\xi_0^{k_1-1}\xi_1\cdots\xi_0^{k_r-1}\xi_1\in S^{10}$ (i.e. $k_1\geq 2$), we put

$$\zeta(w)=\zeta(k_1,\dots,k_r). \tag{29}$$

3.3 Fundamental solutions of KZE1

Proposition 2. The equation KZE1 $\dfrac{dG}{dz}=\Big(\dfrac{X_0}{z}+\dfrac{X_1}{1-z}\Big)G$ has a unique solution $\mathcal{L}(z)$ satisfying the following asymptotic condition;

$$\begin{aligned}&\mathcal{L}(z)=\widehat{\mathcal{L}}(z)z^{X_0},\\ &\widehat{\mathcal{L}}(z) \text{ is holomorphic at } z=0 \text{ and } \widehat{\mathcal{L}}(0)=\mathbf{I}.\end{aligned}$$

Proof. Let us express $\widehat{\mathcal{L}}(z)$ such as $\widehat{\mathcal{L}}(z)=\sum_{s=0}^{\infty}\widehat{\mathcal{L}}_s(z)$ where $\widehat{\mathcal{L}}_s(z)\in\mathcal{U}_s$ is the homogeneous component of the degree s. Then it is easy to show that $\widehat{\mathcal{L}}_s(z)$ satisfies the following recursive relation;

$$\frac{d\widehat{\mathcal{L}}_{s+1}(z)}{dz}=\frac{1}{z}[X_0,\widehat{\mathcal{L}}_s(z)]+\frac{1}{1-z}X_1\widehat{\mathcal{L}}_s(z). \tag{30}$$

Hence we have $\widehat{\mathcal{L}}_0(z)=\mathbf{I}$ and for $s\geq 1$,

$$\begin{aligned}\widehat{\mathcal{L}}_s(z)&=\int_0^z\Big(\xi_0\otimes\mathrm{ad}(X_0)+\xi_1\otimes\mu(X_1)\Big)^s(\mathbf{1}\otimes\mathbf{I})\\ &=\sum_{k_1+\cdots+k_r=s}\mathrm{Li}_{k_1,\dots,k_r}(z)\ \mathrm{ad}(X_0)^{k_1-1}\mu(X_1)\cdots\mathrm{ad}(X_0)^{k_r-1}\mu(X_1)(\mathbf{I}),\end{aligned} \tag{31}$$

where $\mathrm{ad}(X_0)(A)=[X_0,A]$, $\mu(X_1)(A)=X_1A$ for $A\in\mathcal{U}$. Hence $\mathcal{L}(z)$ exists and is unique. □

We call $\mathcal{L}(z)$ a fundamental solution normalized at $z=0$ of (KZE1). We should observe that $\mathcal{L}(z)$ and $\widehat{\mathcal{L}}(z)$ are grouplike elements of $\widetilde{\mathcal{U}}=\mathbf{C}\langle\langle X_0,X_1\rangle\rangle$ (a non-commutative formal power series ring of X_0,X_1), and that $\widehat{\mathcal{L}}(z)$ is single valued and holomorphic in $D=\mathbf{C}\setminus\{z=x\,|\,1\leq x\}$.

Let us find out another representation of $\mathcal{L}(z)$. We set

$$G(z)=\sum_{w\in S}\mathrm{Li}(w;z)W$$

where in the sum of the right hand side, w ranges over the set of words in S, and for $w=\xi_{i_1}\cdots\xi_{i_r}$, we put $W=X_{i_1}\cdots X_{i_r}$. They are dual basis to each other.

We can show the following lemma [HPH], [Ok].

Lemma 3. For any $u \in S$, one has

$$\begin{cases} \dfrac{d}{dz}\,\mathrm{Li}(\xi_0 u; z) = \dfrac{\mathrm{Li}(u;z)}{z}, \\[2ex] \dfrac{d}{dz}\,\mathrm{Li}(\xi_1 u; z) = \dfrac{\mathrm{Li}(u;z)}{1-z}. \end{cases} \tag{32}$$

Hence $G(z)$ is a solution of (KZE1).

Proof. Putting $u = v \mathbin{ш} \xi_0^r$ $(v \in S^0,\ r \geq 0)$, we prove (32) by induction on r. If $r = 0$, then $u = v \in S^0$. This case is already proved because we have

$$\frac{d}{dz}\,\mathrm{Li}_{k_1,k_2,\dots,k_r}(z) = \begin{cases} \dfrac{1}{z}\,\mathrm{Li}_{k_1-1,k_2,\dots,k_r} & (k_1 \geq 2), \\[2ex] \dfrac{1}{1-z}\,\mathrm{Li}_{k_2,\dots,k_r}(z) & (k_1 = 1). \end{cases} \tag{33}$$

Now assume that, for $r-1$ $(r \geq 1)$, (32) holds. Since

$$\xi_i v \mathbin{ш} \xi_0^r = \xi_i(v \mathbin{ш} \xi_0^r) + \xi_0(\xi_i v \mathbin{ш} \xi_0^{r-1}),$$

applying $\mathrm{Li}(\bullet; z)$ to this relation, we have

$$\mathrm{Li}(\xi_i(v \mathbin{ш} \xi_0^r); z) = \mathrm{Li}(\xi_i v; z)\frac{\log^r z}{r!} - \mathrm{Li}(\xi_0(\xi_i v \mathbin{ш} \xi_0^{r-1}); z).$$

By the assumption of induction, the right hand side above reads

$$\begin{aligned} \frac{d}{dz}\{\mathrm{RHS}\} &= \frac{d}{dz}\,\mathrm{Li}(\xi_i v; z)\cdot\frac{\log^r z}{r!} + \frac{1}{z}\,\mathrm{Li}(\xi_i v; z)\frac{\log^{r-1} z}{(r-1)!} - \frac{1}{z}\,\mathrm{Li}(\xi_i v \mathbin{ш} \xi_0^{r-1}; z) \\ &= \frac{d}{dz}\,\mathrm{Li}(\xi_i v; z)\cdot\frac{\log^r z}{r!}. \end{aligned}$$

Since $\xi_i v \in S^0$, we have already proved (32) for $\dfrac{d}{dz}\,\mathrm{Li}(\xi_i v; z)$. Hence we have

$$\frac{d}{dz}\,\mathrm{Li}(\xi_i v; z)\cdot\frac{\log^r z}{r!} = \begin{cases} \dfrac{1}{z}\,\mathrm{Li}(v;z)\dfrac{\log^r z}{r!} = \dfrac{1}{z}\,\mathrm{Li}(v \mathbin{ш} \xi_0^r; z) & (i=0), \\[2ex] \dfrac{1}{1-z}\,\mathrm{Li}(v;z)\dfrac{\log^r z}{r!} = \dfrac{1}{1-z}\,\mathrm{Li}(v \mathbin{ш} \xi_0^r; z) & (i=1). \end{cases}$$

This proves (32) for $u = v \mathbin{ш} \xi_0^r$. □

In [IKZ], the following proposition is implicitly written:

Proposition 4 (IKZ)**.** We put

$$\Phi = \sum_{w \in S} w \otimes W \in S \mathbin{\widehat{\otimes}} \mathcal{U} \tag{34}$$

where $\widehat{\otimes}$ denotes the topological tensor product. Then Φ decomposes as follows:

$$\Phi = \Big(\sum_{w \in S} \mathrm{reg}^0(w) \otimes W\Big) \exp_{ш}(\xi_0 \otimes X_0) \tag{35}$$

where $\exp_{ш}(\xi_0 \otimes X_0) = \sum_{k=0}^{\infty} \xi_0^k \otimes X_0^k$. Furthermore we have

$$\varPhi = \exp_{ш}(\xi_1 \otimes X_1)\Big(\sum_{w \in S} \mathrm{reg}^{10}(w) \otimes W\Big) \exp_{ш}(\xi_0 \otimes X_0). \tag{36}$$

Since, by using (34) and (35), we have

$$G(z) = \big(\mathrm{Li} \otimes \mathrm{id}_{\mathcal{U}}\big)(\varPhi) = \Big(\sum_{w \in S} \mathrm{Li}(\mathrm{reg}^0(w); z) W\Big) z^{X_0},$$

$G(z)$ satisfies the same asymptotic condition as $\mathcal{L}(z)$. Hence

$$G(z) = \mathcal{L}(z).$$

Moreover we obtain the following proposition.

Proposition 5 ([OkU])**.** The fundamental solution $\mathcal{L}(z)$ normalized at $z = 0$ can be written as follows:

$$\begin{aligned} \mathcal{L}(z) &= \sum_{w \in S} \mathrm{Li}(w; z) W && (37) \\ &= \Big(\sum_{w \in S} \mathrm{Li}(\mathrm{reg}^0(w); z) W\Big) z^{X_0} && (38) \\ &= (1 - z)^{-X_1} \Big(\sum_{w \in S} \mathrm{Li}(\mathrm{reg}^{10}(w); z) W\Big) z^{X_0}. && (39) \end{aligned}$$

We consider $\mathcal{L}^{(1)}(z)$, the fundamental solution normalized at $z = 1$ of (KZE1). By the coordinate transformation $z \mapsto t = f(z) = 1 - z$, (KZE1) is transformed to

$$\frac{dG}{dt} = \Big(\frac{-X_1}{t} + \frac{-X_0}{1 - t}\Big) G.$$

Hence we can show that $\mathcal{L}^{(1)}(z)$ which satisfies

$$\mathcal{L}^{(1)}(z) = \widehat{\mathcal{L}}^{(1)}(z)(1 - z)^{-X_1}$$

where $\widehat{\mathcal{L}}^{(1)}(z)$ is holomorphic at $z = 1$ and $\widehat{\mathcal{L}}^{(1)}(1) = \mathbf{I}$, uniquely exists. $\widehat{\mathcal{L}}^{(1)}(z)$ is expressed as follows:

$$\begin{aligned} \widehat{\mathcal{L}}^{(1)}(z) &= \sum_{s=0}^{\infty} \widehat{\mathcal{L}}_s^{(1)}(z), \\ \widehat{\mathcal{L}}_s^{(1)}(z) &= \sum_{k_1 + \cdots + k_r = s} \mathrm{Li}_{k_1, \ldots, k_r}(1 - z) \times \\ &\qquad \times \mathrm{ad}(-X_1)^{k_1 - 1} \mu(-X_0) \cdots \mathrm{ad}(-X_1)^{k_r - 1} \mu(-X_0)(\mathbf{I}). \end{aligned} \tag{40}$$

From Proposition 5, we obtain

$$\begin{aligned} \mathcal{L}^{(1)}(z) &= \sum_{w \in S} \mathrm{Li}(w; 1 - z) f_*(W) && (41) \\ &= \Big(\sum_{w \in S} \mathrm{Li}(\mathrm{reg}^0(w); 1 - z) f_*(W)\Big)(1 - z)^{-X_1} && (42) \\ &= z^{X_0} \Big(\sum_{w \in S} \mathrm{Li}(\mathrm{reg}^{10}(w); 1 - z) f_*(W)\Big)(1 - z)^{-X_1}, && (43) \end{aligned}$$

where $f_* : \mathcal{U} \to \mathcal{U}$ is an automorphism defined by $f_*(X_0) = -X_1$, $f_*(X_1) = -X_0$.

3.4 Drinfel'd associator and the connection problem of KZE1

We set

$$\varPhi_{\mathrm{KZ}} = \varPhi_{\mathrm{KZ}}(X_0, X_1) = \sum_{w \in S} \zeta(\mathrm{reg}^{10}(w))W \tag{44}$$

which is referred to as the Drinfel'd associator. This is also a grouplike element of $\widetilde{\mathcal{U}}$.

Proposition 6 ([OkU])**.** We have

(1) $\mathcal{L}(z) = \mathcal{L}^{(1)}(z)\varPhi_{\mathrm{KZ}}$.

(2) The relation $(\mathcal{L}^{(1)}(z))^{-1}\mathcal{L}(z) = \varPhi_{\mathrm{KZ}}$ is equivalent to

$$\sum_{uv=w} \mathrm{Li}(\tau(u); 1-z)\,\mathrm{Li}(v; z) = \zeta(\mathrm{reg}^{10}(w)) \quad \text{for } \forall w \text{ word in } S. \tag{45}$$

Here $\tau : S \to S$ is an anti-automorphism defined by $\tau(\xi_0) = \xi_1,\ \tau(\xi_1) = \xi_0$.

Proof. (1) Note that $\mathrm{Li}_{k_1,\ldots,k_r}(z)$ is holomorphic in $D = \mathbf{C} \setminus \{z = x \,|\, 1 \le x\}$, and $\mathrm{Li}_{k_1,\ldots,k_r}(1-z)$ is holomorphic in $D' = \mathbf{C} \setminus \{z = x \,|\, x \le 0\}$. From (31) and (40), $\left(\mathcal{L}^{(1)}(z)\right)^{-1}\mathcal{L}(z)$ is holomorphic in a simply connected domain $D \cap D'$. Since $\dfrac{d}{dz}\left\{\left(\mathcal{L}^{(1)}(z)\right)^{-1}\mathcal{L}(z)\right\} = 0$, $\left(\mathcal{L}^{(1)}(z)\right)^{-1}\mathcal{L}(z) = C$ is constant. From (39), (43), taking the limit $z \to 1, z \in D \cap D'$, we have

$$\begin{aligned} C &= \lim_{z \to 1} \left\{(1-z)^{X_1}\Big(\sum_{w \in S} \mathrm{Li}(\mathrm{reg}^{10}(w); 1-z) f_*(W)\Big)^{-1} z^{-X_0} \right. \\ &\qquad \left. \cdot (1-z)^{-X_1}\Big(\sum_{w \in S} \mathrm{Li}(\mathrm{reg}^{10}(w); z)W\Big) z^{X_0}\right\} \\ &= \sum_{w \in S} \zeta(\mathrm{reg}^{10}(w))W = \varPhi_{\mathrm{KZ}} \end{aligned}$$

(2) Since $\mathcal{L}^{(1)}(z)$ is a grouplike element of $\widetilde{\mathcal{U}}$, from (41), we have

$$\left(\mathcal{L}^{(1)}(z)\right)^{-1} = \sum_{w \in S} \mathrm{Li}(w; 1-z)(\rho \circ f_*)(W)$$

where $\rho : \mathcal{U} \to \mathcal{U}$ is an anti-homomorphism defined by $\rho(X_0) = -X_0$, $\rho(X_1) = -X_1$, which is the antipode of $\mathcal{U}$ as a Hopf algebra. Hence, introducing an anti-homomorphism $\tau : S \to S$ defined by $\tau(\xi_0) = \xi_1,\ \tau(\xi_1) = \xi_0$, we have

$$\left(\mathcal{L}^{(1)}(z)\right)^{-1} = \sum_{w \in S} \mathrm{Li}(\tau(w); 1-z)W.$$

Therefore we have

$$\begin{aligned} \left(\mathcal{L}^{(1)}(z)\right)^{-1}\mathcal{L}(z) &= \Big(\sum_{u \in S} \mathrm{Li}(\tau(u); 1-z)U\Big) \cdot \Big(\sum_{v \in S} \mathrm{Li}(v; 1-z)V\Big) \\ &= \sum_{w} \Big(\sum_{uv=w} \mathrm{Li}(\tau(u); 1-z)\,\mathrm{Li}(v; z)\Big)W. \end{aligned}$$

This proves (45).

□

The relation (45) is called the generalized inversion formula.

Proposition 7 ([OkU]). We have

(1) $\Phi_{\mathrm{KZ}}(X_0, X_1)\Phi_{\mathrm{KZ}}(-X_1, -X_0) = \mathbf{I}$ (the duality relation of Φ_{KZ}).

(2) From (1), we obtain the duality relation of MZV:

$$\zeta(\mathrm{reg}^{10}(w)) = \zeta(\mathrm{reg}^{10}(\tau(w))) \quad \text{for } \forall w \in S. \tag{46}$$

Proof. Taking the limit $z \to 0, z \in D \cap D'$ of $(\mathcal{L}^{(1)}(z))^{-1}\mathcal{L}(z)$, we have

$$\Phi_{\mathrm{KZ}}(X_0, X_1) = \Big(\sum_w \mathrm{Li}(\mathrm{reg}^{10}(w); 1) f_*(W)\Big)^{-1} = \Phi_{\mathrm{KZ}}(-X_1, -X_0)^{-1}.$$

Since the Drinfel'd associator is grouplike, we have

$$\begin{aligned}\Phi_{\mathrm{KZ}}(-X_1, -X_0)^{-1} &= \sum_w \zeta(\mathrm{reg}^{10}(w))(\iota \circ f_*)(W) \\ &= \sum_w \zeta(\mathrm{reg}^{10}(\tau(w))W.\end{aligned}$$

This completes the proof. □

For examples of Proposition 6 and Proposition 7, we present the following:

(1) $\mathrm{Li}_2(z) - \log z\, \mathrm{Li}_1(z) + \mathrm{Li}_2(1 - z) = \zeta(2)$. (Euler's inversion formula [Le])

(2) $\zeta(3) = \zeta(2, 1)$. (Indeed, $\zeta(3) = \zeta(\xi_0^2\xi_1) = \zeta(\tau(\xi_0^2\xi_1)) = \zeta(\xi_0\xi_1^2) = \zeta(2, 1)$.)

3.5 Schlesinger type equation of one variable

Consideration on (KZE1) generalizes to the case of the Schlesinger type equation.

$$\frac{dG}{dz} = \Big(\frac{X_0}{z} + \sum_{j=1}^{m} \frac{a_j X_j}{1 - a_j z}\Big)G \tag{SE1}$$

where $a_1, \ldots, a_m \in \mathbf{C} \setminus \{0\}$ are distinct points. The coefficients $X_0, X_1, \ldots, X_m$ are generators of a free Lie algebra

$$\mathfrak{X} = \mathbf{C}\{X_0, X_1, \ldots, X_m\}.$$

The universal enveloping algebra $\mathcal{U} = \mathcal{U}(\mathfrak{X}) = \mathbf{C}\langle X_0, X_1, \ldots, X_m\rangle$ is a non-commutative polynomial ring and a completion $\widetilde{\mathcal{U}} = \widetilde{\mathcal{U}}(\mathfrak{X}) = \mathbf{C}\langle\langle X_0, X_1, \ldots, X_m\rangle\rangle$ is a non-commutative formal power series ring.

The one forms $\xi_0, \xi_1, \ldots, \xi_m$ are defined by

$$\xi_0 = \frac{dz}{z}, \qquad \xi_j = \frac{a_j dz}{1 - a_j z} \quad (1 \le j \le m)$$

and let $S = S(\xi_0, \xi_1, \ldots, \xi_m) = (\mathbf{C}\langle\xi_0, \xi_1, \ldots, \xi_m\rangle, ш, \mathbf{1})$ be a shuffle algebra generated by $\xi_0, \xi_1, \ldots, \xi_m$.

Proposition 8. The Schlesinger type equation (SE1) has a unique solution $\mathcal{L}(z)$ satisfying the asymptotic condition

$$\begin{aligned}&\mathcal{L}(z) = \widehat{\mathcal{L}}(z)z^{X_0},\\ &\widehat{\mathcal{L}}(z) \text{ is holomorphic at } z = 0 \text{ and } \widehat{\mathcal{L}}(0) = \mathbf{I}.\end{aligned}$$

Proof. Similarly as in the case of (KZE1), $\widehat{\mathcal{L}}(z) = \sum_{s=0}^{\infty} \widehat{\mathcal{L}}_s(z)$ ($\widehat{\mathcal{L}}_s(z) \in \mathcal{U}_s$) are expressed as

$$\begin{aligned}\widehat{\mathcal{L}}_s(z) &= \int_0^z \Big(\xi_0 \otimes \mathrm{ad}(X_0) + \sum_{j=1}^{m} \xi_j \otimes \mu(X_j)\Big)^s (\mathbf{1} \otimes \mathbf{I})\\ &= \sum_{\substack{k_1+\cdots+k_r=s\\ i_1,\ldots,i_r\in\{1,\ldots,m\}}} L({}^{k_1}a_{i_1} \cdots {}^{k_r}a_{i_r}; z)\, \mathrm{ad}(X_0)^{k_1-1}\mu(X_{i_1}) \cdots \mathrm{ad}(X_0)^{k_r-1}\mu(X_{i_r})(\mathbf{I}).\end{aligned}$$

Here $L({}^{k_1}a_{i_1} \cdots {}^{k_r}a_{i_r}; z)$ stands for a hyperlogarithm which is by definition

$$\begin{aligned}L({}^{k_1}a_{i_1} \cdots {}^{k_r}a_{i_r}; z) &:= \int_0^z \xi_0^{k_1-1}\xi_{i_1} \cdots \xi_0^{k_r-1}\xi_{i_r}\\ &= \sum_{n_1>n_2\cdots>n_r>0} \frac{a_{i_1}^{n_1-n_2}a_{i_2}^{n_2-n_3} \cdots a_{i_r}^{n_r}}{n_1^{k_1} \cdots n_r^{k_r}} z^{n_1}.\end{aligned}$$

This Taylor expansion is absolutely convergent for $|z| < \min_{1\le k\le r} |a_{i_k}|^{-1}$. □

The solution $\mathcal{L}(z)$ in Proposition 8 is referred to as the fundamental solution normalized at $z = 0$ of (SE1). $\mathcal{L}(z)$, $\widehat{\mathcal{L}}(z)$ are grouplike elements of $\widetilde{\mathcal{U}}$.

4 KZE of two variables

4.1 Definition of KZE of two variables

By the cubic coordinate on $\mathcal{M}_{0,5} : z_1 = r(1, 4; 3, 5), z_2 = r(2, 1; 3, 5)$, KZE on $\mathcal{M}_{0,5}$ is represented as the KZ equation of two variables (KZE2):

$$dG = \varOmega G, \quad \varOmega = \xi_1 X_1 + \xi_{11} X_{11} + \xi_2 X_2 + \xi_{22} X_{22} + \xi_{12} X_{12}, \qquad \text{(KZE2)}$$

$$\xi_1 = \frac{dz_1}{z_1}, \ \ \xi_{11} = \frac{dz_1}{1 - z_1}, \ \ \xi_2 = \frac{dz_2}{z_2}, \ \ \xi_{22} = \frac{dz_2}{1 - z_2}, \ \ \xi_{12} = \frac{d(z_1 z_2)}{1 - z_1 z_2}.$$

The singular divisors D of (KZE2) are

$$D = \{z_1 = 0, 1, \infty\} \cup \{z_2 = 0, 1, \infty\} \cup \{z_1 z_2 = 1\}.$$

The coefficients X_i, X_{ii}, X_{12} are given by

$$\begin{aligned}&X_1 = \varOmega_{12} + \varOmega_{13} + \varOmega_{14}, \quad X_2 = \varOmega_{23},\\ &X_{11} = -\varOmega_{14}, \quad X_{22} = -\varOmega_{12}, \quad X_{12} = -\varOmega_{24},\end{aligned}$$

which satisfy

$$\begin{cases}[X_1, X_2] = [X_{11}, X_2] = [X_1, X_{22}] = 0, \\ [X_{11}, X_{22}] = [-X_{11}, X_{12}] = [X_{22}, X_{12}] = [-X_1 + X_2, X_{12}]. \end{cases} \tag{47}$$

$\mathfrak{X}$ is the Lie algebra generated by $X_1, \dots, X_{12}$ with (47) as the fundamental relations.

Note that (AR) for this case is

$$\begin{cases}\xi_1 \wedge \xi_{11} = 0, \quad \xi_2 \wedge \xi_{22} = 0, \\ (\xi_1 + \xi_2) \wedge \xi_{12} = 0, \\ \xi_{11} \wedge \xi_{12} + \xi_{22} \wedge (\xi_{11} - \xi_{12}) - \xi_2 \wedge \xi_{12} = 0. \end{cases} \tag{48}$$



singular divisors
$D = \{z_1 = 0, 1, \infty\}$
$\cup \{z_2 = 0, 1, \infty\}$
$\cup \{z_1 z_2 = 1\}$

4.2 Fiber space structure of $\mathcal{M}_{0,5}$ and decomposition of $\mathfrak{X}$

Consider the projection

$$p_2 : \mathcal{M}_{0,5} \to \mathcal{M}_{0,4} : \ [x_1, x_2, x_3, x_4, x_5] \mapsto [x_1, x_3, x_4, x_5]. \tag{49}$$



$\mathcal{M}_{0,4} \cong \mathbf{P}^1 \setminus \{0, 1, \infty\}$

From this, we obtain the semi-direct product of the fundamental groups;

$$\pi_1(\mathcal{M}_{0,5}, (z_1, z_2)) \cong \pi_1(\mathbf{P}^1 \setminus \{0, 1, \infty, z_1^{-1}\}, z_2) \rtimes \pi_1(\mathcal{M}_{0,4}, z_1).$$

Taking Lie algebras, we have

$$\mathfrak{X} = \mathbf{C}\{X_2, X_{22}, X_{12}\} \oplus \mathbf{C}\{X_1, X_{11}\}$$

($\mathbf{C}\{X_2, X_{22}, X_{12}\}$ is a Lie ideal). We have similar decomposition

$$\mathfrak{X} = \mathbf{C}\{X_1, X_{11}, X_{12}\} \oplus \mathbf{C}\{X_2, X_{22}\}$$

($\mathbf{C}\{X_1, X_{11}, X_{12}\}$ is a Lie ideal).

Proposition 9. The following decomposition holds:

$$\mathfrak{X} = \mathbf{C}\{X_1, X_{11}, X_{12}\} \oplus \mathbf{C}\{X_2, X_{22}\} = \mathbf{C}\{X_2, X_{22}, X_{12}\} \oplus \mathbf{C}\{X_1, X_{11}\}, \tag{50}$$

$$\begin{aligned} \mathcal{U}(\mathfrak{X}) &= \mathcal{U}(\mathbf{C}\{X_1, X_{11}, X_{12}\}) \otimes \mathcal{U}(\mathbf{C}\{X_2, X_{22}\}) \\ &= \mathcal{U}(\mathbf{C}\{X_2, X_{22}, X_{12}\}) \otimes \mathcal{U}(\mathbf{C}\{X_1, X_{11}\}) \end{aligned} \tag{51}$$

Moreover, $\mathbf{C}\{X_1, X_{11}, X_{12}\}$, $\mathbf{C}\{X_2, X_{22}, X_{12}\}$ are Lie ideals of $\mathfrak{X}$.

4.3 Reduced bar algebra

Now let

$$S = S(\xi_1, \xi_{11}, \xi_2, \xi_{22}, \xi_{12}) = (\mathbf{C}\langle \xi_1, \xi_{11}, \xi_2, \xi_{22}, \xi_{12}\rangle, ш, \mathbf{1})$$

be a shuffle algebra generated by $\xi_1, \dots, \xi_{12}$. We say that an element

$$\varphi = \sum_{I=\{i_1,\dots,i_s\}} c_i \omega_{i_1} \cdots \omega_{i_s}$$

($\omega_{i_k} \in \{\xi_1, \dots, \xi_{12}\}$) satisfies Chen's integrability condition (CIC) if and only if, for $1 \le \forall l \le s-1$,

$$\sum_I c_I \; \omega_{i_1} \otimes \cdots \otimes \omega_{i_l} \wedge \omega_{i_{l+1}} \otimes \cdots \otimes \omega_{i_s} = 0 \tag{CIC}$$

holds as a multiple differential form [C2]. Let $\mathcal{B}$ be a subalgebra of S of elements satisfying (CIC):

$$\mathcal{B} = \{\varphi \in S \mid \varphi \text{ satisfies (CIC)}\}. \tag{52}$$

We refer to $\mathcal{B}$ as the reduced bar algebra.

For $\varphi \in \mathcal{B}$, the iterated integral $\displaystyle\int_{(a_1,a_2)}^{(z_1,z_2)} \varphi$ depends only on the homotopy class of the integral contour and defines a many-valued analytic function on $\mathbf{P}^1 \times \mathbf{P}^1 \setminus D$.

As Hopf algebras, $\mathcal{U}$ and $\mathcal{B}$ are dual to each other.

Proposition 10 ([OU]). There exist isomorphisms of shuffle algebras

$$\iota_{1\otimes 2} : \mathcal{B} \xrightarrow{\cong} S(\xi_1, \xi_{11}, \widetilde{\xi}_{12}^{(1)}) \otimes S(\xi_2, \xi_{22}), \tag{53}$$

$$\iota_{2\otimes 1} : \mathcal{B} \xrightarrow{\cong} S(\xi_2, \xi_{22}, \widetilde{\xi}_{12}^{(2)}) \otimes S(\xi_1, \xi_{11}) \tag{54}$$

where $\widetilde{\xi}_{12}^{(1)} = \dfrac{z_2 dz_1}{1 - z_1 z_2}$, $\widetilde{\xi}_{12}^{(2)} = \dfrac{z_1 dz_2}{1 - z_1 z_2}$ $(\widetilde{\xi}_{12}^{(1)} + \widetilde{\xi}_{12}^{(2)} = \xi_{12})$. The isomorphism $\iota_{1\otimes 2}$ is defined through the following procedure:
(*i*) For $\varphi \in \mathcal{B}$, pick up the terms $\varphi_1\varphi_2 \in S(\xi_1, \xi_{11}, \xi_{12}) \cdot S(\xi_2, \xi_{22})$.
(*ii*) change each term $\varphi_1\varphi_2$ to $\varphi_1 \otimes \varphi_2 \in S(\xi_1, \xi_{11}, \xi_{12}) \otimes S(\xi_2, \xi_{22})$.
(*iii*) replace ξ_{12} to $\widetilde{\xi}_{12}^{(1)}$.
$\iota_{2\otimes 1}$ is similarly defined.

Let $\mathcal{B}^0$ be a subalgebra of elements ending with other than ξ_1, ξ_2:

$$\mathcal{B}^0 = \{\varphi \in \mathcal{B} \mid \varphi = \sum c_I \omega_{i_1} \cdots \omega_{i_s},\ \omega_{i_s} \neq \xi_1, \xi_2\}.$$

For $\varphi \in \mathcal{B}^0$, we can put an iterated integral $\displaystyle\int_{(0,0)}^{(z_1,z_2)} \varphi$.

Proposition 11 ([OU])**.** We have isomorphisms of shuffle algebras

$$\iota_{1\otimes 2} : \mathcal{B}^0 \xrightarrow{\cong} S^0(\xi_1, \xi_{11}, \widetilde{\xi}^{(1)}_{12}) \otimes S^0(\xi_2, \xi_{22}), \tag{55}$$

$$\iota_{2\otimes 1} : \mathcal{B}^0 \xrightarrow{\cong} S^0(\xi_2, \xi_{22}, \widetilde{\xi}^{(2)}_{12}) \otimes S^0(\xi_1, \xi_{11}), \tag{56}$$

where $S^0(A) = \{\varphi \in S(A) \mid \varphi = \sum c_I \omega_{i_1} \cdots \omega_{i_s},\ \omega_{i_s} \neq \xi_1, \xi_2\}$.

4.4 Fundamental solution of KZE2

Let $\Omega_0 = \xi_1 X_1 + \xi_2 X_2$ (the singular part of Ω at $(0,0)$) and $\Omega' = \Omega - \Omega_0$ (the regular part of Ω at $(0,0)$).

Proposition 12. KZE2 has a unique solution $\mathcal{L}(z_1, z_2)$ satisfying the asymptotic condition:

$$\begin{aligned}&\mathcal{L}(z_1, z_2) = \widehat{\mathcal{L}}(z_1, z_2) z_1^{X_1} z_2^{X_2},\\ &\widehat{\mathcal{L}}(z_1, z_2) \text{ is holomorphic at } (0,0) \text{ and } \widehat{\mathcal{L}}(0,0) = \mathbf{I}.\end{aligned}$$

Proof. Since the fundamental relations (47) of $\mathfrak{X}$ is homogeneous, $\mathcal{U}$ has the canonical gradation: $\mathcal{U} = \displaystyle\bigoplus_{s=0}^{\infty} \mathcal{U}_s$. Similarly, the reduced bar algebra $\mathcal{B}$ and the subalgebra $\mathcal{B}^0$ are graded: $\mathcal{B} = \displaystyle\bigoplus_{s=0}^{\infty} \mathcal{B}_s,\ \mathcal{B}^0 = \bigoplus_{s=0}^{\infty} \mathcal{B}^0_s \quad (B^0_s = B_s \cap B^0)$.

Let $\widehat{\mathcal{L}}(z_1, z_2) = \displaystyle\sum_{s=0}^{\infty} \widehat{\mathcal{L}}_s(z_1, z_2),\ (\widehat{\mathcal{L}}_s \in \mathcal{U}_s)$. Then we have the following recursive relation:

$$d\widehat{\mathcal{L}}_{s+1}(z_1, z_2) = [\Omega_0, \widehat{\mathcal{L}}_s] + \Omega' \widehat{\mathcal{L}}_s.$$

Lemma 13 ([OU], Proposition 17)**.** We have

$$\left(\mathrm{ad}(\Omega_0) + \mu(\Omega')\right)^s (\mathbf{1} \otimes \mathbf{I}) \in \mathcal{B}^0_s \otimes \mathcal{U}_s$$

where we set $\mathrm{ad}(\omega \otimes X)(\varphi \otimes A) = \omega\varphi \otimes [X, A]$, $\mu(\omega \otimes X)(\varphi \otimes A) = \omega\varphi \otimes XA$.

Hence we have $\widehat{\mathcal{L}}_0(z) = \mathbf{I}$, and for $s \geq 1$

$$\widehat{\mathcal{L}}_s(z_1, z_2) = \int_{(0,0)}^{(z_1,z_2)} \left(\mathrm{ad}(\Omega_0) + \mu(\Omega')\right)^s (\mathbf{1} \otimes \mathbf{I}). \tag{57}$$

This shows that the fundamental solution $\mathcal{L}(z_1, z_2)$ exists and is unique. □

The solution $\mathcal{L}(z_1, z_2)$ is called the fundamental solution normalized at $(0,0)$ of (KZE2) . We should observe that $\mathcal{L}(z_1, z_2)$ and $\widehat{\mathcal{L}}(z_1, z_2)$ are grouplike elements of $\widetilde{\mathcal{U}}$ which is the completion of $\mathcal{U}$ with respect to the gradation, and that $\widehat{\mathcal{L}}(z_1, z_2)$ is single valued and holomorphic in $\{(z_1, z_2) \,|\, |z_1| < 1,\ |z_2| < 1\}$.

4.5 Decomposition theorem of fundamental solution

Proposition 14. The fundamental solution $\mathcal{L}(z_1, z_2)$ normalized at $(0, 0)$ decomposes as

$$\mathcal{L}(z_1, z_2) = \mathcal{L}^{(2)}_{2\otimes 1}(z_1, z_2)\mathcal{L}^{(1)}_{2\otimes 1}(z_1) = \mathcal{L}^{(1)}_{1\otimes 2}(z_1, z_2)\mathcal{L}^{(2)}_{1\otimes 2}(z_2). \tag{$\sharp$}$$

Here $\mathcal{L}^{(j)}_{i\otimes j}(z_j)$ is the fundamental solution normalized at $z_j = 0$ of

$$\frac{dG}{dz_j} = \left(\frac{X_j}{z_j} + \frac{X_{jj}}{1 - z_j}\right) G \tag{KZE1$_{i\otimes j}$}$$

$$\mathcal{L}^{(j)}_{i\otimes j}(z_j) = \widehat{\mathcal{L}}^{(j)}_{i\otimes j}(z_j) z_j^{X_j}, \quad \widehat{\mathcal{L}}^{(j)}_{i\otimes j}(0) = \mathbf{I}.$$

$\mathcal{L}^{(i)}_{i\otimes j}(z_1, z_2)$ is the fundamental solution normalized at $z_i = 0$ of

$$\frac{dG}{dz_i} = \left(\frac{X_i}{z_i} + \frac{X_{ii}}{1 - z_i} + \frac{z_j X_{12}}{1 - z_1 z_2}\right) G \tag{SE1$_{i\otimes j}$}$$

$$\mathcal{L}^{(i)}_{i\otimes j}(z_1, z_2) = \widehat{\mathcal{L}}^{(i)}_{i\otimes j}(z_1, z_2) z_i^{X_i}, \quad \widehat{\mathcal{L}}^{(i)}_{i\otimes j}\Big|_{z_i=0} = \mathbf{I}.$$

The holomorphic part $\widehat{\mathcal{L}}(z_1, z_2)$ has similar decomposition as follows:

$$\widehat{\mathcal{L}}(z_1, z_2) = \widehat{\mathcal{L}}^{(2)}_{2\otimes 1}(z_1, z_2)\widehat{\mathcal{L}}^{(1)}_{2\otimes 1}(z_1) = \widehat{\mathcal{L}}^{(1)}_{1\otimes 2}(z_1, z_2)\widehat{\mathcal{L}}^{(2)}_{1\otimes 2}(z_2). \tag{$\sharp'$}$$

Proof. From the asymptotic condition of $\mathcal{L}(z_1, z_2) = \widehat{\mathcal{L}}(z_1, z_2) z_1^{X_1} z_2^{X_2}$, it follows that $G(z_1) = \widehat{\mathcal{L}}(z_1, 0) z_1^{X_1}$ is the fundamental solution of (KZE1$_{2\otimes 1}$) . Hence $G(z_1) = \mathcal{L}^{(1)}_{2\otimes 1}(z_1)$. Put

$$H(z_1, z_2) = \mathcal{L}(z_1, z_2)\big(\mathcal{L}^{(1)}_{2\otimes 1}(z_1)\big)^{-1}.$$

Then $H(z_1, z_2)$ is a solution of (SE1$_{2\otimes 1}$). Since $[X_2, X_1] = [X_2, X_{11}] = 0$, it has the asymptotic condition as follows:

$$\begin{aligned} &H(z_1, z_2) = \hat{H}(z_1, z_2) z_2^{X_2}, \\ &\widehat{H}(z_1, z_2) = \widehat{\mathcal{L}}(z_1, z_2)\big(\widehat{\mathcal{L}}(z_1, 0)\big)^{-1}, \quad \widehat{H}(z_1, 0) = \mathbf{I}. \end{aligned}$$

Therefore $H(z_1, z_2) = \mathcal{L}^{(2)}_{2\otimes 1}(z_1, z_2)$. □

We call this proposition “Decomposition theorem for the fundamental solution”.

5 Connection Problem of KZE on $\mathcal{M}_{0,5}$

5.1 Compactification of $\mathcal{M}_{0,5}$

A smooth compactification of $\mathcal{M}_{0,5}$ is given as follows [Y]:

$$\overline{\mathcal{M}}_{0,5} = \mathrm{PGL}(2, \mathbf{C})\Big\backslash\big((\mathbf{P}^1)^5 \setminus \Delta'\big), \tag{58}$$

$$\Delta' = \{(x_1, x_2, x_3, x_4, x_5) \in (\mathbf{P}^1)^5 \mid \text{at least 3 points coincide}\}.$$

Then

$$\overline{\mathcal{M}}_{0,5} \setminus \mathcal{M}_{0,5} = \bigcup_{1 \le i < j \le 5} D_{ij}, \tag{59}$$

$$D_{ij} = \{[x_1, x_2, x_3, x_4, x_5] \mid x_i = x_j\}. \tag{60}$$

The divisors D_{ij}'s are rational curves in $\overline{\mathcal{M}}_{0,5}$ and satisfy

$$D_{ij} \cap D_{kl} = \begin{cases} \emptyset & (\{i,j\} \cap \{k,l\} \neq \emptyset,\ \{i,j\} \neq \{k,l\}), \\ \{\text{one point}\} & (\{i,j\} \cap \{k,l\} = \emptyset). \end{cases}$$

Let $\overline{\mathcal{M}}_{0,5}(\mathbf{R})$ be the real points of $\overline{\mathcal{M}}_{0,5}$:

$$\mathcal{M}_{0,5}(\mathbf{R}) = \overline{\mathcal{M}}_{0,5}(\mathbf{R}) \setminus \bigcup_{1 \le i < j \le 5} D_{ij}(\mathbf{R}).$$

Then $\mathcal{M}_{0,5}(\mathbf{R})$ has 12 connected components, and each component is a pentagon surrounded by five curves $D_{ij}(\mathbf{R})$:



In the above figure, pentagons with the same number should be identified.

5.2 Cubic coordinate system associated with a pentagon

Let us fix the pentagon 1:



Each vertex of this pentagon is given as follows:

$$P_0=[0,1,1,\infty,\infty],\quad P_1=[0,\infty,\infty,1,0],\quad P_2=[\infty,0,1,\infty,1],$$
$$P_3=[1,1,\infty,0,\infty],\quad P_4=[\infty,\infty,0,1,1].$$

The cubic coordinate system

$$\begin{aligned}(z_1,z_2)&=\big(r(1,4;3.5),\ r(2,1;3.5)\big)\\ &=\left(\frac{(x_1-x_3)(x_4-x_5)}{(x_1-x_5)(x_4-x_3)},\frac{(x_2-x_3)(x_1-x_5)}{(x_2-x_5)(x_1-x_3)}\right)\end{aligned}\tag{61}$$

is a local coordinate system around P_0, and via the map

$$[x_1,x_2,x_3,x_4,x_5]\ \mapsto\ (z_1,z_2)$$

the pentagon 1 blows down to the square $\{(z_1,z_2)\,|\,0\le z_1\le 1,\ 0\le z_2\le 1\}$ as follows:

$$(z_1,z_2)(P_0)=(0,0),\ (z_1,z_2)(D_{23})=\{z_2=0\},\ (z_1,z_2)(D_{45})=\{z_1=0\},$$
$$(z_1,z_2)(D_{14})=\{z_1=1\},\ (z_1,z_2)(D_{12})=\{z_2=1\},\ (z_1,z_2)(D_{35})=(1,1).$$

We should observe that $\mathcal{L}(z_1,z_2)$, the fundamental solution normalized at $(z_1,z_2)=(0,0)$ of (KZE2), is a fundamental solution of (KZE) which is single valued and holomorphic in a certain domain containing the pentagon 1.

Put $\sigma=\begin{pmatrix}1&2&3&4&5\\5&4&1&3&2\end{pmatrix}$. Define $\sigma:\overline{\mathcal{M}}_{0,5}\to\overline{\mathcal{M}}_{0,5}$ as follows: For a point $P=[a_1,a_2,a_3,a_4,a_5]$, we set

$$\begin{aligned}\sigma(P)&=[a_{\sigma^{-1}(1)},a_{\sigma^{-1}(2)},a_{\sigma^{-1}(3)},a_{\sigma^{-1}(4)},a_{\sigma^{-1}(5)}]\\ &\quad(=[a_3,a_5,a_4,a_2,a_1]).\end{aligned}\tag{62}$$

Then σ is an automorphism (attached to the pentagon 1) of $\overline{\mathcal{M}}_{0,5}$ satisfying $\sigma^5=\mathrm{id}$ and

$$\sigma(P_\alpha)=P_{\alpha+1}\ (\mathrm{mod}\ 5),\quad \sigma(D_{ij})=D_{\sigma(i),\sigma(j)}\quad(\forall i,j).\tag{63}$$

Let $\sigma^*(x_i)=x_{\sigma(i)}$ $(i=1,2,3,4,5)$ which induces an automorphism of the rational function field of $\overline{\mathcal{M}}_{0,5}$. Put $(z_1^{(0)},z_2^{(0)})=(z_1,z_2)$. Then

$$\begin{cases}z_1^{(1)}=\sigma^*(z_1^{(0)})=r(5,3;1,2)=z_2^{(0)},\\[2ex] z_2^{(1)}=\sigma^*(z_2^{(0)})=r(4,5;1,2)=\dfrac{1-z_1^{(0)}}{1-z_1^{(0)}z_2^{(0)}}\end{cases}$$

are local coordinates around P_1. In general, we set

$$\begin{cases} z_1^{(\alpha+1)} = \sigma^*(z_1^{(\alpha)}) = z_2^{(\alpha)}, \\ \\ z_2^{(\alpha+1)} = \sigma^*(z_2^{(\alpha)}) = \dfrac{1-z_1^{(\alpha)}}{1-z_1^{(\alpha)}z_2^{(\alpha)}}. \end{cases} \tag{64}$$

Then $(z_1^{(\alpha)}, z_2^{(\alpha)})$ is a local coordinate system around P_α, and is called the cubic coordinate system attached to P_α.

Define the pull-back of ω_{ij} by

$$\sigma^*(\omega_{ij}) = d\log(\sigma^*(x_i) - \sigma^*(x_j)), \tag{65}$$

and let $\sigma_* : \mathfrak{X} \to \mathfrak{X}$ be an induced automorphism defined through

$$\sum_{1\le i<j\le 5} \omega_{ij}\sigma_*(\Omega_{ij}) = \sum_{1\le i<j\le 5} \sigma^*(\omega_{ij})\Omega_{ij}, \tag{66}$$

namely,

$$\sigma_*(\Omega_{ij}) = \Omega_{\sigma^{-1}(i),\sigma^{-1}(j)}. \tag{67}$$

Then we have

$$\Omega = (\sigma^* \otimes \sigma_*^{-1})(\Omega) = \sum_{1\le i<j\le 5} \sigma^*(\omega_{ij})\sigma_*^{-1}(\Omega_{ij}). \tag{68}$$

We represent Ω by the cubic coordinate system attached to P_α: Let

$$\begin{aligned} &X_i^{(0)} = X_i, \quad X_{ij}^{(0)} = X_{ij} \quad (1 \le i \le 2,\ 1 \le i \le j \le 2), \\ &X_i^{(\alpha+1)} = \sigma_*^{-1}(X_i^{(\alpha)}), \quad X_{ij}^{(\alpha+1)} = \sigma_*^{-1}(X_{ij}^{(\alpha)}) \quad (\alpha = 0,1,2,3). \end{aligned}$$

Then Ω reads

$$\Omega = \xi_1^{(\alpha)}X_1^{(\alpha)} + \xi_{11}^{(\alpha)}X_{11}^{(\alpha)} + \xi_2^{(\alpha)}X_2^{(\alpha)} + \xi_{22}^{(\alpha)}X_{22}^{(\alpha)} + \xi_{12}^{(\alpha)}X_{12}^{(\alpha)}, \tag{69}$$

where

$$\xi_i^{(\alpha)} = \frac{dz_i^{(\alpha)}}{z_i^{(\alpha)}}, \quad \xi_{ii}^{(\alpha)} = \frac{dz_i^{(\alpha)}}{1-z_i^{(\alpha)}} \quad (i=1,2), \quad \xi_{12}^{(\alpha)} = \frac{d(z_1^{(\alpha)}z_2^{(\alpha)})}{1-z_1^{(\alpha)}z_2^{(\alpha)}}.$$

Let $\mathcal{L}^{(\alpha)} = \mathcal{L}^{(\alpha)}(z_1^{(\alpha)}, z_2^{(\alpha)})$ be the fundamental solution normalized at P_α of KZE2$_\alpha$

$$dG = \left(\xi_1^{(\alpha)}X_1^{(\alpha)} + \xi_{11}^{(\alpha)}X_{11}^{(\alpha)} + \xi_2^{(\alpha)}X_2^{(\alpha)} + \xi_{22}^{(\alpha)}X_{22}^{(\alpha)} + \xi_{12}^{(\alpha)}X_{12}^{(\alpha)}\right)G. \tag{KZE2$_\alpha$}$$

We should observe that each $\mathcal{L}^{(\alpha)}$ is a fundamental solution of (KZE) which is single valued and holomorphic in a certain domain containing the pentagon 1.

Let us find the connection matrix C_α;

$$\mathcal{L}^{(\alpha)}(z_1^{(\alpha)}, z_2^{(\alpha)}) = \mathcal{L}^{(\alpha+1)}(z_1^{(\alpha+1)}, z_2^{(\alpha+1)})C_\alpha. \tag{70}$$

To this equation, we substitute the decomposition (♯) for $\mathcal{L}^{(\alpha)}, \mathcal{L}^{(\alpha+1)}$

$$\mathcal{L}^{(\alpha)}(z_1^{(\alpha)}, z_2^{(\alpha)}) = \mathcal{L}_{2\otimes 1}^{(\alpha)(2)}(z_1^{(\alpha)}, z_2^{(\alpha)})\mathcal{L}_{2\otimes 1}^{(\alpha)(1)}(z_1^{(\alpha)}),$$
$$\mathcal{L}^{(\alpha+1)}(z_1^{(\alpha+1)}, z_2^{(\alpha+1)}) = \mathcal{L}_{1\otimes 2}^{(\alpha+1)(1)}(z_1^{(\alpha+1)}, z_2^{(\alpha+1)})\mathcal{L}_{1\otimes 2}^{(\alpha+1)(2)}(z_2^{(\alpha+1)}),$$

where $\mathcal{L}_{i\otimes j}^{(\beta)(j)}(z_j^{(\beta)})$ $(\beta = \alpha, \alpha+1)$ is the fundamental solution normalized at $z_j^{(\beta)} = 0$ of $\mathrm{KZE1}_{i\otimes j}^{(\beta)}$

$$\frac{dG}{dz_j^{(\beta)}} = \left(\frac{X_j^{(\beta)}}{z_j^{(\beta)}} + \frac{X_{jj}^{(\beta)}}{1 - z_j^{(\beta)}}\right) G \qquad (\mathrm{KZE1}_{i\otimes j}^{(\beta)})$$
$$\mathcal{L}_{i\otimes j}^{(\beta)(j)}(z_j) = \widehat{\mathcal{L}}_{i\otimes j}^{(\beta)(j)}(z_j^{(\beta)})\bigl(z_j^{(\beta)}\bigr)^{X_j^{(\beta)}}, \quad \widehat{\mathcal{L}}_{i\otimes j}^{(\beta)(j)}(0) = \mathbf{I},$$

and $\mathcal{L}_{i\otimes j}^{(\beta)(i)}(z_1^{(\beta)}, z_2^{(\beta)})$ is the fundamental solution normalized at $z_i^{(\beta)} = 0$ of $\mathrm{SE1}_{i\otimes j}^{(\beta)}$

$$\frac{dG}{dz_i^{(\beta)}} = \left(\frac{X_i^{(\beta)}}{z_i^{(\beta)}} + \frac{X_{ii}^{(\beta)}}{1 - z_i^{(\beta)}} + \frac{z_j^{(\beta)}X_{12}^{(\beta)}}{1 - z_1^{(\beta)}z_2^{(\beta)}}\right) G \qquad (\mathrm{SE1}_{i\otimes j}^{(\beta)})$$
$$\mathcal{L}_{i\otimes j}^{(\beta)(i)}(z_1^{(\beta)}, z_2^{(\beta)}) = \widehat{\mathcal{L}}_{i\otimes j}^{(\beta)(i)}(z_1^{(\beta)}, z_2^{(\beta)})\bigl(z_i^{(\beta)}\bigr)^{X_i^{(\beta)}}, \quad \widehat{\mathcal{L}}_{i\otimes j}^{(\beta)(i)}\Big|_{z_i^{(\beta)}=0} = \mathbf{I}.$$

Then we have

$$\begin{aligned}\mathcal{L}_{2\otimes 1}^{(\alpha)(2)}(z_1^{(\alpha)})C_\alpha^{-1} = \mathcal{L}_{2\otimes 1}^{(\alpha)(2)}(z_1^{(\alpha)}, z_2^{(\alpha)})^{-1}\mathcal{L}_{1\otimes 2}^{(\alpha+1)(1)}(z_1^{(\alpha+1)}, z_2^{(\alpha+1)})& \\ \cdot\, \mathcal{L}_{1\otimes 2}^{(\alpha+1)(2)}(z_2^{(\alpha+1)}).&\end{aligned} \qquad (71)$$

Note that $z_1^{(\alpha+1)} = z_2^{(\alpha)}$, $z_2^{(\alpha+1)} = \dfrac{1 - z_1^{(\alpha)}}{1 - z_1^{(\alpha)}z_2^{(\alpha)}}$ and $X_1^{(\alpha+1)} = X_2^{(\alpha)}$, $X_2^{(\alpha+1)} = -X_{11}^{(\alpha)}$. Since the left hand side of (71) is independent of the variable $z_2^{(\alpha)}$, first taking the limit of $z_2^{(\alpha)} \to 0$, we have

$$\begin{aligned}\mathrm{RHS} &= \mathcal{L}_{1\otimes 2}^{(\alpha+1)(2)}(1 - z_1^{(\alpha)})\bigl(1 - z_1^{(\alpha)}\bigr)^{-X_{11}^{(\alpha)}} \\ &\sim \mathbf{I} \times (1 - z_1^{(\alpha)})^{-X_{11}^{(\alpha)}} \quad (z_1^{(\alpha)} \to 1).\end{aligned}$$

Hence RHS is the fundamental solution normalized at $z_1^{(\alpha)} = 0$ of $\mathrm{KZE1}_{i\otimes j}^{(\alpha)}$. This implies $C_\alpha = \varPhi_{\mathrm{KZ}}(X_1^{(\alpha)}, X_{11}^{(\alpha)})$.

Theorem 15. Put $\varPhi_{\mathrm{KZ}}^{(\alpha)} = \varPhi_{\mathrm{KZ}}(X_1^{(\alpha)}, X_{11}^{(\alpha)})$.

(1) We have

$$\mathcal{L}^{(\alpha)} = \mathcal{L}^{(\alpha+1)}\varPhi_{\mathrm{KZ}}^{(\alpha)} \quad (\alpha = 0, 1, 2, 3, 4 \pmod 5)). \qquad (72)$$

(2) As the compatibility condition for the connection relation (72), we obtain the pentagon relation of the Drinfel'd associator

$$\varPhi_{\mathrm{KZ}}^{(4)}\,\varPhi_{\mathrm{KZ}}^{(3)}\,\varPhi_{\mathrm{KZ}}^{(2)}\,\varPhi_{\mathrm{KZ}}^{(1)}\,\varPhi_{\mathrm{KZ}}^{(0)} = \mathbf{I}. \qquad (73)$$

Proof. We show (2). Since the connection relations (72) hold in a certain domain containing the pentagon 1, we have

$$\mathcal{L}^{(0)} = \mathcal{L}^{(1)}\varPhi_{\mathrm{KZ}}^{(0)} = \mathcal{L}^{(2)}\varPhi_{\mathrm{KZ}}^{(1)}\varPhi_{\mathrm{KZ}}^{(0)} = \cdots = \mathcal{L}^{(0)}\varPhi_{\mathrm{KZ}}^{(4)}\varPhi_{\mathrm{KZ}}^{(3)}\varPhi_{\mathrm{KZ}}^{(2)}\varPhi_{\mathrm{KZ}}^{(1)}\varPhi_{\mathrm{KZ}}^{(0)}.$$

From this we obtain the pentagon relation (73). □

6 Five term relation of dilogarithms

6.1 Iterated integral representation along $C_{1\otimes 2},\ C_{2\otimes 1}$ of $\mathcal{L}(z_1, z_2)$

Let $C_{1\otimes 2},\ C_{2\otimes 1}$ be contours defined in the following figure:



For $\varphi_2 \otimes \varphi_1 \in S^0(\xi_2, \xi_{22}, \widetilde{\xi}_{12}^{(2)}) \otimes S^0(\xi_1, \xi_{11})$, we set

$$\int_{2\otimes 1} \varphi_2 \otimes \varphi_1 := \int_{C_{2\otimes 1}} \varphi_2\varphi_1 = \int_{z_2=0}^{z_2} \varphi_2 \int_{z_1=0}^{z_1} \varphi_1. \tag{74}$$

Similarly for $\psi_1 \otimes \psi_2 \in S^0(\xi_1, \xi_{11}, \widetilde{\xi}_{12}^{(1)}) \otimes S^0(\xi_2, \xi_{22})$, we set

$$\int_{1\otimes 2} \psi_1 \otimes \psi_2 := \int_{C_{1\otimes 2}} \psi_1\psi_2 = \int_{z_1=0}^{z_1} \psi_1 \int_{z_2=0}^{z_2} \psi_2. \tag{75}$$

Proposition 16. For $\varphi \in \mathcal{B}^0$, we have

$$\begin{aligned}\int_{(0,0)}^{(z_1,z_2)} \varphi = \int_{C_{2\otimes 1}} \varphi &= \int_{2\otimes 1} \iota_{2\otimes 1}(\varphi) \\ = \int_{C_{1\otimes 2}} \varphi &= \int_{1\otimes 2} \iota_{1\otimes 2}(\varphi).\end{aligned}$$

Hence we have

$$\int_{2\otimes 1} \iota_{2\otimes 1}(\varphi) = \int_{1\otimes 2} \iota_{1\otimes 2}(\varphi). \tag{76}$$

We call (76) a generalized harmonic product relation. This provides a relation of hyperlogarithms of the type $\mathcal{M}_{0.5}$ which is defined as follows: For $\alpha_i \in \{1, z_2\} \quad (i = 1, \ldots, r)$, we set

$$\begin{aligned} L({}^{k_1}\alpha_1 \cdots {}^{k_r}\alpha_r; z_1) :&= \int_0^{z_1} \xi_1^{k_1-1}\omega_1 \cdots \xi_1^{k_r-1}\omega_r \\ &= \sum_{n_1 > \cdots > n_r > 0} \frac{\alpha_1^{n_1-n_2}\alpha_2^{n_2-n_3}\cdots\alpha_r^{n_r}}{n_1^{k_1}\cdots n_r^{k_r}} z_1^{n_1}, \end{aligned} \tag{77}$$

where $\omega_i = \dfrac{\alpha_i dz_1}{1-\alpha_i z_1}$. This is called hyperlogarithms of the type $\mathcal{M}_{0.5}$ of the main variable z_1. If $\alpha_1 = \cdots = \alpha_r = 1$, it reduces to a MPL of one variable;

$$\mathrm{Li}_{k_1,\ldots,k_r}(z_1) = L({}^{k_1}1 \cdots {}^{k_r}1\,; z_1). \tag{78}$$

If $\alpha_1 = \cdots = \alpha_i = 1,\ \alpha_{i+1} = \cdots = \alpha_{i+j} = z_2\ (r = i + j)$, it is a MPL of two variables;

$$\mathrm{Li}_{k_1,\ldots,k_r}(i, r-i; z_1, z_2) = \sum_{n_1 > n_2 \cdots > n_r > 0} \frac{z_1^{n_1} z_2^{n_{i+1}}}{n_1^{k_1}\cdots n_r^{k_r}}. \tag{79}$$

The generalized harmonic product relations are the main subject in [OU]. Here we only give a simple example.

Let $\varphi = \xi_{11}\xi_{12} + \xi_{22}\xi_{11} - \xi_{22}\xi_{12} - \xi_2\xi_{12} \in \mathcal{B}^0$. Then

$$\begin{aligned} &\iota_{1\otimes 2}(\varphi) = \xi_{11}\widetilde{\xi}_{12}^{(1)} \otimes \mathbf{1}, \\ &\iota_{2\otimes 1}(\varphi) = \xi_{22} \otimes \xi_{11} - \xi_{22}\widetilde{\xi}_{12}^{(2)} \otimes \mathbf{1} - \xi_2\widetilde{\xi}_{12}^{(2)} \otimes \mathbf{1}, \end{aligned}$$

$$\begin{aligned} \int_{1\otimes 2} \iota_{1\otimes 2}(\varphi) &= \int_0^{z_1} \xi_{11}\widetilde{\xi}_{12}^{(1)} = \mathrm{Li}_{1,1}(1,1; z_1, z_2), \\ \int_{2\otimes 1} \iota_{2\otimes 1}(\varphi) &= \int_0^{z_2} \xi_{22} \int_0^{z_1} \xi_{11} - \int_0^{z_2} \xi_{22}\widetilde{\xi}_{12}^{(2)} - \int_0^{z_2} \xi_2\widetilde{\xi}_{12}^{(2)} \\ &= \mathrm{Li}_1(z_2)\,\mathrm{Li}_1(z_1) - \mathrm{Li}_{1,1}(1,1; z_2, z_1) - \mathrm{Li}_2(0,1 : z_2, z_1). \end{aligned}$$

Thus (76) for φ reads

$$\mathrm{Li}_1(z_1)\,\mathrm{Li}_1(z_2) = \mathrm{Li}_{1,1}(1,1; z_1, z_2) + \mathrm{Li}_2(0,1; z_2, z_1) + \mathrm{Li}_{1,1}(1,1; z_2, z_1). \tag{80}$$

From Proposition 16, we have

$$\begin{aligned} \widehat{\mathcal{L}}_s(z_1, z_2) &= \int_{C_{1\otimes 2}} (\mathrm{ad}(\Omega_0) + \mu(\Omega'))^s\,(\mathbf{1} \otimes \mathbf{I}) \\ &= \int_{C_{1\otimes 2}} (\iota_{1\otimes 2} \otimes \mathrm{id}_{\mathcal{U}})\left((\mathrm{ad}(\Omega_0) + \mu(\Omega'))^s\,(\mathbf{1} \otimes \mathbf{I})\right), \end{aligned}$$

where $\widehat{\mathcal{L}}_s(z_1, z_2)$ is the homogeneous degree s part of the fundamental solution $\mathcal{L}(z_1, z_2)$ of KZE2. Hence we have the following proposition.

Proposition 17 ([OU], Corollary 21)**.** (1) We have

$$\widehat{\mathcal{L}}_s(z_1, z_2) = \sum_{s'+s''=s} \sum_{W',W''} \int_0^{z_1} \theta^{(1)}_{1\otimes 2}(W') \int_0^{z_2} \theta^{(2)}_{1\otimes 2}(W'') \ \alpha(W')\alpha(W'')(\mathbf{I}) \tag{81}$$

where W' runs over the set $\mathcal{W}^0_{s'}(X_1, X_{11}, X_{12})$, and W'' runs over the set $\mathcal{W}^0_{s''}(X_2, X_{22})$. ($\mathcal{W}^0_s(\mathfrak{A}) = \mathcal{W}^0(\mathfrak{A}) \cap \mathcal{U}_s(\mathfrak{X})$, and $\mathcal{W}^0(\mathfrak{A})$ stands for the set of words of the letters $\mathfrak{A}$ ending with other than X_1, X_2.) $\alpha : \mathcal{U}(\mathfrak{X}) \to \mathrm{End}(\mathcal{U}(\mathfrak{X}))$ is an algebra homomorphism

$$\alpha : (X_1, X_{11}, X_2, X_{22}, X_{12}) \mapsto (\mathrm{ad}(X_1), \mu(X_{11}), \mathrm{ad}(X_2), \mu(X_{22}), \mu(X_{12})),$$

and

$$\begin{aligned} \theta^{(1)}_{1\otimes 2} &: \mathcal{U}(\mathbf{C}\{X_1, X_{11}, X_{12}\}) \to S(\xi_1, \xi_{11}, \widetilde{\xi}^{(1)}_{12}) \\ \theta^{(2)}_{1\otimes 2} &: \mathcal{U}(\mathbf{C}\{X_2, X_{22}\}) \to S(\xi_2, \xi_{22}) \end{aligned}$$

are linear maps defined by replacing

$$\theta^{(i)}_{1\otimes 2}(X_i) = \xi_i, \ \theta^{(i)}_{1\otimes 2}(X_{ii}) = \xi_{ii} \quad (i = 1, 2), \ \theta^{(1)}_{1\otimes 2}(X_{12}) = \widetilde{\xi}^{(1)}_{12}.$$

(2) Similarly we have

$$\widehat{\mathcal{L}}_s(z_1, z_2) = \sum_{s'+s''=s} \sum_{W',W''} \int_0^{z_2} \theta^{(2)}_{2\otimes 1}(W') \int_0^{z_1} \theta^{(1)}_{2\otimes 1}(W'') \ \alpha(W')\alpha(W'')(\mathbf{I}) \tag{82}$$

where W' runs over the set $\mathcal{W}^0_{s'}(X_2, X_{22}, X_{12})$, and W'' runs over the set $\mathcal{W}^0_{s''}(X_1, X_{11})$, and

$$\begin{aligned} \theta^{(2)}_{2\otimes 1} &: \mathcal{U}(\mathbf{C}\{(X_2, X_{22}, X_{12}\}) \to S(\xi_2, \xi_{22}, \widetilde{\xi}^{(2)}_{12}) \\ \theta^{(1)}_{2\otimes 1} &: \mathcal{U}(\mathbf{C}\{X_1, X_{11}\}) \to S(\xi_1, \xi_{11}) \end{aligned}$$

are linear maps defined by replacing

$$\theta^{(i)}_{2\otimes 1}(X_i) = \xi_i, \ \theta^{(i)}_{2\otimes 1}(X_{ii}) = \xi_{ii} \quad (i = 1, 2), \ \theta^{(2)}_{2\otimes 1}(X_{12}) = \widetilde{\xi}^{(2)}_{12}.$$

In this proposition, we should observe that the iterated integral

$$L\big(\theta^{(1)}_{1\otimes 2}(W')\,;\, z_1\big) := \int_0^{z_1} \theta^{(1)}_{1\otimes 2}(W') \quad (W' \in \mathcal{W}^0_{s'}(X_1, X_{11}, X_{12}))$$

is a hyperlogarithm of the type $\mathcal{M}_{0,5}$ of the main variable z_1, and the iterated integral

$$L\big(\theta^{(2)}_{1\otimes 2}(W'')\,;\, z_2\big) := \int_0^{z_2} \theta^{(2)}_{1\otimes 2}(W'') \quad (W'' \in \mathcal{W}^0_{s''}(X_2, X_{22}))$$

is a multiple polylogarithm of the variable z_2.

6.2 Five term relation of dilogarithms

Let $\tau = (23)(45)$. Define an automorphism $\tau : \overline{\mathcal{M}}_{0,5} \to \overline{\mathcal{M}}_{0,5}$ as follows: For a point $P = [a_1, a_2, a_3, a_4, a_5]$, set

$$\tau(P) = [a_{\tau^{-1}(1)}, a_{\tau^{-1}(2)}, a_{\tau^{-1}(3)}, a_{\tau^{-1}(4)}, a_{\tau^{-1}(5)}]. \tag{83}$$

By this automorphism, the pentagon 1 is transformed to the pentagon 4 like as in the following figure.



Let $\tau^*(x_i) = x_{\tau(i)}$ $(i = 1, 2, 3, 4, 5)$, which induces an automorphism of the rational function field of $\overline{\mathcal{M}}_{0,5}$, in particular, the cubic coordinates are transformed as follows:

$$\tau^*(z_1, z_2) = \left(\frac{-z_1(1-z_2)}{1-z_1}, \frac{-z_2(1-z_1)}{1-z_2} \right). \tag{84}$$

Define the pull-back of ω_{ij} by $\tau^*(\omega_{ij}) = d\log(\tau^*(x_i) - \tau^*(x_j))$, and let $\tau_* : \mathfrak{X} \to \mathfrak{X}$ be an automorphism defined by

$$\sum_{1 \le i < j \le 5} \omega_{ij} \tau_*(\mathit{\Omega}_{ij}) = \sum_{1 \le i < j \le 5} \tau^*(\omega_{ij}) \mathit{\Omega}_{ij},$$

namely, $\tau_*(\mathit{\Omega}_{ij}) = \mathit{\Omega}_{\tau^{-1}(i), \tau^{-1}(j)}$. Then we have

$$\mathit{\Omega} = (\tau^* \otimes \tau_*^{-1})(\mathit{\Omega}) = \sum_{1 \le i < j \le 5} \tau^*(\omega_{ij}) \tau_*^{-1}(\mathit{\Omega}_{ij}).$$

Explicit representations of the pull-back τ^* and the induced automorphism τ_* are given as follows:

$$\begin{cases} \tau^*(\xi_1) = \xi_1 + \xi_{11} - \xi_{22}, \quad \tau^*(\xi_1) = -\xi_1 + \xi_{12}, \\ \tau^*(\xi_2) = -\xi_{11} + \xi_2 + \xi_{22}, \quad \tau^*(\xi_{22}) = -\xi_{22} + \xi_{12}, \quad \tau^*(\xi_{12}) = \xi_{12}, \end{cases} \tag{85}$$

and

$$\begin{cases} \tau_*(X_1) = X_1, \quad \tau_*(X_{11}) = X_1 - X_{11} - X_2, \quad \tau_*(X_2) = X_2 \\ \tau_*(X_{22}) = -X_1 + X_2 - X_{22}, \quad \tau_*(X_{12}) = X_{11} + X_{22} + X_{12}. \end{cases} \tag{86}$$

Since $\Omega = (\tau^* \otimes \tau_*^{-1})(\Omega)$,

$$\begin{aligned}\widetilde{\mathcal{L}}(z_1, z_2) &= (\tau^* \otimes \tau_*^{-1})(\mathcal{L}(z_1, z_2)) \\ &= \mathcal{L}(\tau^*(z_1), \tau^*(z_2))\Big|_{X \to \tau_*^{-1} X,\ \ X = X_1, X_{11}, X_2, X_{22}, X_{12}}\end{aligned}$$

is also a fundamental solution of (KZE2) which has the asymptotic condition

$$\widetilde{\mathcal{L}}(z_1, z_2) \sim \mathbf{I} \left(\frac{-z_1(1-z_2)}{1-z_1} \right)^{X_1} \left(\frac{-z_2(1-z_1)}{1-z_2} \right)^{X_2} \qquad (z_1, z_2) \to (0, 0). \tag{87}$$

Therefore the connection relation of $\mathcal{L}(z_1, z_2)$ and $\widetilde{\mathcal{L}}(z_1, z_2)$ is

$$\widetilde{\mathcal{L}}(z_1, z_2) = \mathcal{L}(z_1, z_2) \exp(-\operatorname{sgn}(\operatorname{Im} z_1)\, \pi i X_1)\, \exp(-\operatorname{sgn}(\operatorname{Im} z_2)\, \pi i X_2). \tag{88}$$

It is convenient to rewrite this formula as follows:

$$\begin{aligned}(\tau^* \otimes \mathrm{id})(\mathcal{L}(z_1, z_2)) &= (\mathrm{id} \otimes \tau_*^{-1})(\mathcal{L}(z_1, z_2)) \\ &\quad \times \exp(-\operatorname{sgn}(\operatorname{Im} z_1)\, \pi i X_1) \exp(-\operatorname{sgn}(\operatorname{Im} z_2)\, \pi i X_2).\end{aligned} \tag{89}$$

Using (81) and (82) in Proposition 17, we have the following proposition.

Proposition 18. (1) From the coefficients of $[X_1, X_{11}]$ of the both sides of (89), we have

$$\begin{aligned}\mathrm{Li}_2\left(\frac{-z_1(1-z_2)}{1-z_1} \right) &= \mathrm{Li}_{1,1}(1, 1; z_1, z_2) - \mathrm{Li}_2(z_1) - \mathrm{Li}_{1,1}(z_1) \\ &\quad + \mathrm{Li}_2(0, 1; z_1, z_2).\end{aligned} \tag{L1}$$

(2) From the coefficients of $[X_2, X_{22}]$ of both sides of (89), we have

$$\begin{aligned}\mathrm{Li}_2\left(\frac{-z_2(1-z_1)}{1-z_2} \right) &= -\mathrm{Li}_{1,1}(1, 1; z_1, z_2) - \mathrm{Li}_2(z_2) - \mathrm{Li}_{1,1}(z_2) \\ &\quad + \mathrm{Li}_1(z_2)\, \mathrm{Li}_1(z_1).\end{aligned} \tag{L2}$$

The formula (L1) is a two dimensional analogue of Landen's formula of dilogarithms [Le]:

$$\mathrm{Li}_2\left(\frac{-z}{1-z} \right) = -\mathrm{Li}_2(z) - \frac{1}{2} \log^2(1-z).$$

Since $\mathrm{Li}_2(0, 1; z_1, z_2) = \mathrm{Li}_2(z_1 z_2)$ and $\mathrm{Li}_{1,1}(z_1) = \frac{1}{2} \log^2(1-z_1)$, (L1) + (L2) implies the following:

Theorem 19. We have the five term relation for dilogarithms [Le]:

$$\begin{aligned}\mathrm{Li}_2(z_1 z_2) = \mathrm{Li}_2\left(\frac{-z_1(1-z_2)}{1-z_1} \right) + \mathrm{Li}_2\left(\frac{-z_2(1-z_1)}{1-z_2} \right) + \mathrm{Li}_2(z_1) + \mathrm{Li}_2(z_2) \\ + \frac{1}{2} \log^2\left(\frac{1-z_1}{1-z_2} \right).\end{aligned} \tag{5TR}$$

References

[A] V.I.Arnold, The cohomology ring of the colored braid group, Mat. Zametki **5** (1969), pp. 227-231: Math. Notes **5** (1969), pp. 138-140.

[B] F.Brown, Multiple zeta values and periods of moduli spaces $\overline{\mathfrak{M}}_{0,n}(\mathbf{R})$, Ann. Sci. Éc. Norm. Supér. (4) 42 (2009), no. 3, 371-489.

[C1] K.T.Chen, Algebras of iterated path integrals and fundamental groups, Transaction of the American Math. Soc. **156** (1971), 359–379.

[C2] K.T.Chen, Reduced bar constructions on de Rham complexes, Algebra, topology, and category theory (a collection of papers in honor of Samuel Eilenberg), pp. 19–32. Academic Press, New York, 1976.

[Dr] V.G.Drinfel'd, On quasitriangular quasi-Hopf algebras and on a group that is closely connected with Gal($\overline{\mathbf{Q}}/\mathbf{Q}$), Algebra i Analiz 2 (1990), no. 4, 149–181; translation in Leningrad Math. J. **2** (1991), no. 4, 829–860.

[Ha] R.M.Hain, On a generalization of Hilbert's 21st problem, Ann. scient.Éc. Norm. Sup., 4^e série, t. 19, (1986), 609–627.

[HPH] Hoang Ngoc Minh, M. Petitot, and J. van der Hoeven, L'algèbre des polylogarithmes par les séries génératrices. Proc. of FPSAC'99, 11-th International Conference of Formal Power Series and Algebraic Combinatrics, Barcelona, June 1999.

[I] Y.Ihara, Automorphisms of pure sphere braid groups and Galois representations, The Grothendieck Festschrift, Vol. II, 353–373, Progr. Math., 87, Birkhäuser Boston, Boston, MA, 1990.

[IKZ] K.Ihara, M.Kaneko and D.Zagier, Derivation and double shuffle relations for multiple zeta values, Compos. Math. 142 (2006), no. 2, 307–338.

[Le] L.Lewin, The Evolution of the Ladder Concept, Structural Properties of Polylogarithms (ed. by L.Lewin), Math. Surveys and Monographs, **37**, Amer. MAth. Soc. (1991),1–23.

[Ok] J.Okuda, Duality Formulas of Special Values of Multiple Polylogarithms, Bull. London Math. Soc. (2004)

[OkU] J.Okuda and K.Ueno, The Sum Formula for Multiple Zeta Values, Zeta Functions, Topology and Quantum Physics, Developments in Mathematics 14, ed. by T. Aoki et al., Springer-Verlag (2005), 145-170.

[OU] S.Oi and K.Ueno, KZ equation on the moduli space $\mathcal{M}_{0,5}$ and the harmonic product of multiple polylogarithms, preprint (2011) arXiv:math.QA/0910.0718v3.

[R] C.Reutenauer, Free Lie Algebras, Oxford Science Publications,1993.

[Y] M.Yoshida, Hypergeomtric Functions, My Love, Aspects of Mathematics, E 32, Vieweg, 1997.

[W] Z.Wojtkowiak, Monodromy of iterated integrals and non-abelian unipotent periods, Geometric Galois actions, 2, 219–289, London Math. Soc. Lecture Note Ser., 243, Cambridge Univ. Press, Cambridge, 1997.